

RESEARCH ARTICLE

Open Access

# Heterologous hybridisation to a *Pinus* microarray: profiling of gene expression in *Pinus radiata* saplings exposed to ethephon

Joseph G Dubouzet<sup>1</sup>, Lloyd Donaldson<sup>1</sup>, Michael A Black<sup>2</sup>, Les McNoe<sup>2</sup>, Vincent Liu<sup>1</sup> and Gareth Lloyd-Jones<sup>1\*</sup>

## Abstract

**Background:** The design, construction and application of a *Pinus* microarray platform are described. The oligonucleotide microarray was developed using publicly available *Pinus* cDNA sequences mostly derived from *Pinus taeda* to test whether heterologous hybridisation of microarray probes will generate useful data when hybridised with cRNA constructed from the dominant New Zealand forestry species *Pinus radiata*.

**Methods:** A comprehensive consensus sequence collection of *Pinus* cDNA sequences was collated into a non-redundant database used for automated design of 60-mer oligonucleotide microarray probes. The microarray slides, manufactured by Agilent Technologies (Palo Alto, California), were used to monitor gene expression in an induction experiment using 2-chloroethylphosphonic acid, common name ethephon and the active ingredient of the plant growth regulator Ethrel® (Bayer Crop Science). The transcriptomes from tissues of 2-year old *Pinus radiata* saplings +/- ethephon treatment were compared by hybridisation onto the *Pinus* microarray slides.

**Results:** Statistically significant differentially expressed genes identified by heterologous hybridisation to the *Pinus* microarray following ethephon induction included the up-regulation of genes in the xylem that were related to the metabolism of phenylpropanoids and flavonoids, and also defence responses, specifically against fungal/insect attack and oxidative stress. Bark, mucilaginous xylem and xylem generated largely mutually exclusive cohorts of genes and Gene Ontology (GO) classes. The results are also interpreted in reference to gross and microscopic morphological changes. Samples of gene responses were validated by quantitative RT-PCR.

**Conclusion:** These results confirm the successful development of a *Pinus* microarray and demonstrate the utility of the microarray for transcriptomic research in *Pinus radiata* through heterologous hybridisation.

**Keywords:** Heterologous hybridisation; Microarray; *Pinus*; Ethephon; Ethylene

## Background

Microarray technology has been extensively used to profile gene expression for more than a decade. The ability to perform up to a million parallel hybridisations on a single microarray provides researchers with a very efficient platform for characterisation of genetic processes in target organisms. In addition to characterising the transcriptomic differences generated from an experiment involving one species, the same microarray platform can be used for research involving related species through heterologous hybridisation (Davey et al. 2009). Microarrays can also be

used for studies involving methylation and epigenetics by hybridisation with bisulfite-treated genomic DNA (Reinders et al. 2008). This versatility in other downstream applications helps to ensure that microarray technology continues to be relevant in crop research, despite the availability and widely recognised superiority of RNA-Seq for transcriptomic analyses.

*Pinus radiata* D. Don is grown in 1.6 M ha in New Zealand, making it the premier species for silviculture (New Zealand Forest Owners Association 2011). Microarrays based on comprehensive transcriptomic data can be used to advance pine research, however most examples of *Pinus* microarrays published in the last 10 years are based on very modest collections containing less

\* Correspondence: gareth.lloyd-jones@scionresearch.com

<sup>1</sup>Scion, 49 Sala St., Private Bag 3020, Rotorua, New Zealand

Full list of author information is available at the end of the article

than 3,500 sequenced genes (Heath et al. 2002; Yang et al. 2004; Myburg et al. 2006; Adomas et al. 2007; Balk et al. 2007; Perdiguero et al. 2012; Li et al. 2012). A cDNA microarray based on 26,496 sequences for *P. taeda* was recently used to assess the effects of drought in *P. taeda* L. (Lorenz et al. 2011).

As of July 2013, there were approximately 13,000 publicly available nucleotide sequences from *P. radiata* curated at GenBank (<http://www.ncbi.nlm.nih.gov/>), but much larger collections of DNA sequences exist from other *Pinus* species, principally *P. taeda*, which had approximately 140,000 sequences. Microarrays can be used for comparative genomic hybridisation among near relatives via heterologous hybridisation (Davey et al. 2009). Heterologous hybridisation between cDNA probes from *P. taeda*, immobilised on nylon membrane, with cDNA from *P. sylvestris* L. and *Picea abies* (L.) H. Karst has been successful (Van Zyl et al. 2002). Heterologous hybridisation requires microarray probes that are capable of tolerating a certain degree of mismatch with the target transcripts. Single nucleotide deletions in targets of 60-mer probes can reduce signal intensity by 20-50% whereas single deletions in targets of 20-mer probes essentially eliminated any signal (LeProust 2008). Probes of 60-mer can still generate useful signals even with 10% mismatched bases between the probe and target gene (Hughes et al. 2001).

The utility of a microarray is best appreciated by the degree to which the results from this technology confirm or clarify well-characterised biological phenomena. The effect of long-term exposure to an ethylene-generating paste containing 2-chloroethylphosphonic acid (ethephon) on the stem anatomy of *Pinus radiata* saplings was used as a physiological model to test heterologous hybridisation to the *Pinus* microarray. The response of xylogenesis to ethylene has not previously been studied adequately at the molecular level in *Pinus* species. Ethephon breaks down in plant tissues to form ethylene, a major plant hormone with diverse effects on the plant's physiology (Warner and Leopold 1969). It is known that different *Pinus* species show clear responses to ethephon treatment. Previously, the application of an ethylene-generating paste has been found to induce localised radial stem growth in white pine (*Pinus strobus* L.) (Brown and Leopold 1973). Ethylene also stimulated growth in bark and xylem tissues of *Pinus halepensis* Mill. (Yamamoto and Kozlowski 1987a). Ethephon reduced lignification of tracheids in compression wood that developed on the underside of tilted stems of *Pinus densiflora* Siebold & Zucc. (Yamamoto and Kozlowski 1987b). In *P. halepensis*, ethephon application implicated ethylene in the control of resin duct formation (Yamamoto and Kozlowski 1987a). Ethephon sprays of 1–2 g L<sup>-1</sup> effectively reduced height and radial growth of jack pine (*Pinus banksiana* Lamb.) (Schnurr et al. 1996).

The central assumption of this study is that heterologous microarray hybridisation can be used to generate useful data with probes from closely related species. Microarray probes were designed primarily from *P. taeda* hybridised with cRNA constructed from the close relative *P. radiata*. The creation of a comprehensive consensus sequence database for *Pinus* species, the design and construction of a microarray platform from this database, and the use of this microarray platform to monitor and characterise the transcriptomic response of *P. radiata* saplings to ethephon will be described. The results of heterologous hybridisation were verified by comparing signal data reported from specific probes in the array with physiological observations and, for a limited number of probes, by quantitative reverse transcription polymerase chain reaction (RT-PCR). The lists of differentially expressed genes were analysed using the Database for Annotation, Visualisation and Integrated Discovery (DAVID) Bioinformatics Resources (Huang et al. 2008) and are discussed in relation to gross and microscopic changes induced by ethephon application.

## Methods

### Plant materials

One-year-old rooted clonal saplings of a single genotype of *Pinus radiata* were purchased from ArborGen Australasia (Whakatane, New Zealand) for this experiment. These were placed in 500 mL pots containing standard potting mix that were then arrayed on an unroofed concrete pavement for another year prior to their use in this research. A total of 60 saplings were required to provide 15 plants per treatment for 4 treatments.

### Ethephon application

Lanolin pastes (0.2 to 0.5 g) containing 0, 1, 5 and 25 mg of ethephon supplied from ETHREL (48% Ethephon, Bayer Crop Science, NZ) were dispensed onto 3 × 6 cm aluminium foil sheets that were wrapped around the basal stem of each experimental sapling, 5–7.5 cm from the soil level. The 1 mg per plant dosage corresponded to the 1% in lanolin dose used in *P. halepensis* (Yamamoto and Kozlowski 1987a).

### RNA extraction


RNA was extracted from bark, mucilaginous xylem (containing the cambium) and xylem samples. Bark refers to the outermost layer of tissues of the stem beyond the mucilaginous developing xylem. Xylem refers to the woody fibrous tissue left after scraping off the mucilaginous developing xylem. RNA was extracted from the 0 and 1 mg treatments for transcriptomic analysis. The 5 and 25 mg ethephon treatments at 8 weeks after treatment (WAT) elicited the greatest physiological responses

but transcriptomic analyses were not attempted for these treatments due to inconsistent RNA extraction possibly due to the adverse effect of high resin content. Three replicates consisting of 5 saplings per replicate were destructively sampled 1 week after ethephon treatment and an identical batch was sampled 8 weeks after treatment. Thus each tissue was sampled from each of 15 saplings for a given treatment. After wiping off the residual lanolin from the bark surface with tissue paper, the bark was cut and peeled off, the mucilaginous xylem was scraped off with the blunt edge of a box cutter, and the remaining woody xylem was cut off from the rest of the stem. The respective tissues were wrapped in waterproof paper, flash-frozen in liquid nitrogen and stored at  $-80^{\circ}\text{C}$  prior to RNA extraction. Total RNA from xylem was extracted using Concert reagent (Invitrogen; Carlsbad, CA) whilst RNA extracted from bark was obtained using Concert reagent followed by purification with the Plant RNA Isolation kit (Agilent). RNA from the mucilaginous xylem was extracted using the Plant RNA Isolation kit.

#### Assembly and annotation of the heterologous *Pinus* Consensus Transcript database

In mid-2010, there were 18,900 Unigene entries for *P. taeda* in the US National Center for Biotechnology Information (NCBI) database ([ftp://ftp.ncbi.nih.gov/repository/UniGene/Pinus\\_taeda/Pta.seq.uniq.gz](ftp://ftp.ncbi.nih.gov/repository/UniGene/Pinus_taeda/Pta.seq.uniq.gz)). The Plant Genome Database (<http://www.plantgdb.org/>) had 77,500 Putative Unique Transcripts (PUT), ([http://www.plantgdb.org/download/download.php?dir=/Sequence/ESTcontig/Pinus\\_taeda/current\\_version/Pinus\\_taeda.mRNA.PUT.fasta.bz2](http://www.plantgdb.org/download/download.php?dir=/Sequence/ESTcontig/Pinus_taeda/current_version/Pinus_taeda.mRNA.PUT.fasta.bz2)). Harvard's Computational Biology and Functional Genomics Laboratory (<http://compbio.dfci.harvard.edu/cgi-bin/tgi/gimain.pl?gudb=pine>) had 39,800 tentative consensus (TC) sequences ([ftp://occams.dfci.harvard.edu/pub/bio/tgi/data/Pinus/PGL.release\\_8.zip](ftp://occams.dfci.harvard.edu/pub/bio/tgi/data/Pinus/PGL.release_8.zip)). These sequences were combined into a FASTA file generated using BioEdit software (Hall 1999) and then reassembled using the default parameters for sequence assembly using CAP3 software ([http://seq.cs.iastate.edu/CAP3/cap3.linux.i686\\_xeon64.tar](http://seq.cs.iastate.edu/CAP3/cap3.linux.i686_xeon64.tar)) (Huang and Madan 1999). The final database consisted of 68,225 sequences, with 39,869 assembled contigs in common between at least two of the three databases and 28,356 'singletons' unique to one database. The assembled sequences ranged from 50 to 6175 base pairs (bp), with an average length of 811 bp. The distribution of the assembled sequences is shown in Figure 1. The reassembled database, henceforth referred to as the All-*Pinus* database, was used for the design of 60-mer probes for the *Pinus* microarray.

The most similar protein for each sequence assembly was determined using the BLASTX algorithm as implemented with the BioEdit software, using the default



criteria and a maximum E-value set at 0.001. The closest protein was identified as the one with the most significant E-value among those identified by BLASTX. The actual average E-value for the whole database was  $1.9 \times 10^{-5}$ . With these criteria, 40,238 (59%) sequences were assigned a putative protein ID, of which 14,838 were unique. A cut-off E-value set at  $1 \times 10^{-5}$  was used in the annotation of Conifer DBMagic (Lorenz et al. 2012) and a smaller number of putative protein IDs (36,995 sequences (54%)) within the All-*Pinus* database sequences were identified when this cut-off E-value was used. The annotations for these protein IDs were downloaded from DAVID v6.7, (<http://david.abcc.ncifcrf.gov/>) (Huang et al. 2009). The top 40 sequences identified by comparison with the Swiss-Prot database of annotated protein sequences and the numbers of transcript variants that code for essentially the same protein are shown in Table 1.

#### Pre-testing the All-*Pinus* sequence database

The 7,538 expressed sequence tags (ESTs) from *P. radiata* found in the US National Center for Biotechnology Information (NCBI) database as of 2010 were used to query the 68,225 sequences in the All-*Pinus* database via BLASTN algorithm (Altschul et al. 1997). The BLASTN algorithm was set at a cut-off value of  $1.0 \times E^{-10}$ . The alignments that generated less than the highest score for each EST were eliminated. The average data from BLASTN (630 bp aligned length, 99.2% identity, 619 bp actual length, 12.7 unaligned bases, 3.7 mismatched bases and 0.8 gaps) reinforced the potential of heterologous hybridisation of *P. radiata* sequences to a *Pinus* microarray developed from the All-*Pinus* database.

#### Microarray Probe design and *in silico* validation

Candidate probes were selected based on base-composition profiles optimised through eArray (<https://earray.chem>).

**Table 1 Top 40 most common proteins identified in the All-Pinus database**

SwissProt ID	Description and original species	Number of sequences	Proportion
Q75JC9	Uncharacterised protein DDB_G0271670 OS = <i>Dictyostelium discoideum</i>	33	0.20%
O94317	Uncharacterised serine-rich protein C215.13 OS = <i>Schisosaccharomyces pombe</i>	27	0.16%
Q17RH7	Putative protein TPRXL OS = <i>Homo sapiens</i>	26	0.16%
P26986	2S seed storage-like protein OS = <i>Picea glauca</i>	19	0.12%
Q8LGG8	Universal stress protein A-like protein OS = <i>Arabidopsis thaliana</i>	18	0.11%
P37219	Abscisic stress-ripening protein 2 OS = <i>Solanum lycopersicum</i>	15	0.09%
P19037	18.1 kDa class I heat shock protein OS = <i>Arabidopsis thaliana</i>	14	0.09%
O65351	Subtilisin-like protease OS = <i>Arabidopsis thaliana</i>	13	0.08%
Q6J163	Auxin-induced protein SNG4 OS = <i>Pinus taeda</i> PE = 2 SV = 1	13	0.08%
P13240	Disease resistance reprotein 206 OS = <i>Pisum sativum</i>	12	0.07%
Q00451	36.4 kDa proline-rich protein OS = <i>Solanum lycopersicum</i>	12	0.07%
Q40392	TMV resistance protein N OS = <i>Nicotiana glutinosa</i>	12	0.07%
Q50EK4	Cytochrome P450 750A1 OS = <i>Pinus taeda</i>	11	0.07%
Q8VZ13	Uncharacterised protein At1g08160 OS = <i>Arabidopsis thaliana</i>	11	0.07%
Q9S7C9	Putative DNA-binding protein ESCAROLA OS = <i>Arabidopsis thaliana</i>	11	0.07%
C0LGT6	LRR receptor-like serine/threonine-protein kinase EFR OS = <i>Arabidopsis thaliana</i>	10	0.06%
O80337	Ethylene-retranscription factor 1A OS = <i>Arabidopsis thaliana</i>	9	0.06%
P27061	Acid pho1 OS = <i>Solanum lycopersicum</i>	9	0.06%
Q766C3	Aproteainase nepenthesin-1 OS = <i>Nepenthes gracilis</i>	9	0.06%
Q8BT18	Serine/arginine repetitive matrix protein 2 OS = <i>Mus musculus</i>	9	0.06%
O23530	Protein SUPPRESSOR OF npr1-1, CONSTITUTIVE 1 OS = <i>Arabidopsis thaliana</i>	8	0.05%
P17407	21 kDa protein OS = <i>Daucus carota</i>	8	0.05%
P50160	Sex determination protein tasselseed-2 OS = <i>Zea mays</i>	8	0.05%
Q02817	Mucin-2 OS = <i>Homo sapiens</i>	8	0.05%
Q39224	Protein SRG1 OS = <i>Arabidopsis thaliana</i>	8	0.05%
Q41001	Blue copper protein OS = <i>Pisum sativum</i>	8	0.05%
Q50EK3	Cytochrome P450 704C1 OS = <i>Pinus taeda</i>	8	0.05%
Q55909	TVP38/TMEM64 family membrane protein slr0305 OS = <i>Synechocystis</i>	8	0.05%
Q5ZJC8	mTERF domain-containing protein 1, mitochondrial OS = <i>Gallus gallus</i>	8	0.05%
Q766C2	Aproteainase nepenthesin-2 OS = <i>Nepenthes gracilis</i>	8	0.05%
Q84UV8	Bifunctional monodehydroascorbate reductase and carbonic anhydrase nectarin-3 OS = <i>Nicotiana langsdorffii</i> x <i>Nicotiana sanderae</i>	8	0.05%
Q95JC9	Basic proline-rich protein OS = <i>Sus scrofa</i>	8	0.05%
Q9LHE3	Protein APROTEASE IN GUARD CELL 2 OS = <i>Arabidopsis thaliana</i>	8	0.05%
Q9SB61	ZF-HD homeobox protein At4g24660 OS = <i>Arabidopsis thaliana</i>	8	0.05%
Q9SEV0	Anthocyanidin reductase OS = <i>Arabidopsis thaliana</i>	8	0.05%
Q9SJL0	UDP-glycosyltransferase 86A1 OS = <i>Arabidopsis thaliana</i>	8	0.05%
Q9SZN7	Heavy metal-associated isoprenylated plant protein 26 OS = <i>Arabidopsis thaliana</i>	8	0.05%
Q9UQ35	Serine/arginine repetitive matrix protein 2 OS = <i>Homo sapiens</i>	8	0.05%
Q9ZNU2	NAC domain-containing protein 18 OS = <i>Arabidopsis thaliana</i>	8	0.05%
P05790	Fibroin heavy chain OS = <i>Bombyx mori</i>	7	0.04%

agilent.com/earray/). Three 60-mer sense probes were selected across the length of the template sequence. The 191,500 candidate probes were trimmed to 175,614 to fit

the 180 k format required by the Agilent software. Probes were eliminated based on high similarity and hybridisation quality as predicted by eArray. The remaining probes were

used to query the original All-*Pinus* database by BLASTN, which was set at a cut-off value of  $1.0 \times E^{-10}$ . The results indicated that 123,311 probes (70.2%) for 49,081 genes (71.9%) would give a target-specific signal. The remaining probes were expected to hybridise to sequences such as paralogues and splice variants that have different accession numbers but are essentially different versions of the same protein. The probe sequences were used to query the 7538 ESTs from *P. radiata* in the GenBank sequence database provided by the NCBI. The average identity, aligned length, non-identical bases and gaps were 99.6%, 59.9 bp, 0.2 bp and 0.0 bp respectively. These results affirmed the potential utility of the All-*Pinus* database for generating heterologous *Pinus* microarray probes for analysing transcriptomes derived from *P. radiata*.

#### Microarray analysis

Paired hybridisation for each microarray comprised treatment versus control of each tissue from the same time point, for example; 'bark, 0 mg ethephon, 1 WAT' versus 'bark, 1 mg ethephon, 1 WAT', and so forth. Total RNA samples of 50–100 ng were used for labelling with the fluorescent cyanine dyes Cy3 or Cy5 using the Agilent Low input Quick Amp labelling kit. After labelling, unincorporated dye was removed using RNeasy (Qiagen, Valencia, CA) mini-spin columns. Purified samples were assessed for yield and specific activity using a Nanodrop ND-1000 UV–VIS spectrophotometer (Thermo Fisher Scientific, Wilmington, DE). Samples that gave yields of  $>0.8 \mu\text{g}$  cRNA and specific activity (Concentration of Cy3/Concentration of cRNA  $\times 1000$ )  $>10$  were used for arrays. Labelled samples were hybridised to arrays at 65°C for 17 hours at 10 rpm in an Agilent G2545A hybridisation oven using 2 $\times$  GEx Agilent hybridisation buffer. Slides were washed using Agilent wash buffers. Array scanning and feature extraction was performed using the Agilent G2565CA microarray scanner as outlined in the Agilent protocol for Two-colour Microarray-Based Gene Expression Analysis, Version 6.5, May 2010.

Data were imported into R software using the limma package (Smyth 2005), transformed to the log scale (base 2), and normalized via loess smoothing with no background correction. Statistical analysis of data was performed using the linear models approach implemented in limma, using the M values for each array as the dependent data. The empirical Bayes shrinkage-based approach to per-probe variance estimation was used to generate a moderated t-statistic for each probe (Smyth 2005). This statistic was then used to determine the statistical significance of the differences for each gene at each time/tissue combination. Probability (*P*) values were calculated for each comparison, and used to define the False Discovery Rate threshold by the

Benjamin-Hochberg method (Benjamini and Hochberg 1995).

Probes with adjusted *P*-values greater than 0.01, log<sub>2</sub> fold changes between –1 and 1, and probes with no Uniprot homologs were eliminated. The remaining significantly expressed probes, divided into up- and down-regulated subgroups, were uploaded onto the DAVID portal (<http://david.abcc.ncifcrf.gov/home.jsp>) for singular enrichment analysis in comparison with the list of protein sequences in the All-*Pinus* database. The number of protein IDs associated with each gene ontology classification that were significantly different at  $p < 0.01$  and displayed a conservative False Discovery Rate less than 10% are shown in Table 2.

#### Quantitative RT-PCR verification of microarray signals

Ten probes were selected that generated stable signals greater than background across all treatment combinations after heterologous hybridisation of the *Pinus* microarray; the probes included Contig12730c, which showed strong stable signals that did not change with ethephon/sampling time for use as a control. Forward and reverse primers were designed to amplify  $>45$  bp fragments within the 60-mer probe using OligoAnalyzer 3.1 (<http://www.idtdna.com/analyzer/Applications/OligoAnalyzer/>). Primer details are shown in Table 3 for the successful primer pairs. Aliquots from the original samples that contained 200 to 1300 ng RNA were reverse transcribed using qScript (Quanta Biosciences, MD, USA) in 20  $\mu\text{l}$  reactions. A 1:20 dilution was added as template for a qPCR mix using Lightcycler® Faststart DNA Masterplus SYBR Green i (Roche, USA). A Lightcycler® Carousel-based system (Roche, USA) was used for thermal cycling. A 10 min denaturation step was followed by 45 thermal cycles consisting of a 10 sec denaturation step at 95°C, annealing step of 4 sec at 60°C and an extension step of 3 sec at 72°C. This was followed by melting point analysis from 65°C to 95°C at a ramp rate of 0.2°C sec<sup>-1</sup>. Fold-change calculation, using the expression level of Contig12730c as the constant divisor, was performed with a formula (3.5) for relative quantification (Pfaffl 2004).

#### Microscopy

The gross physiological effects of ethephon were analysed by microscopic analysis of sectioned stems of all treatments. Stem cross-sections were prepared with a sledge microtome and images of the cut ends of stems were obtained using a stereo microscope. Stems were fixed in FAA (5% formalin, 5% glacial acetic acid, 90% ethanol v/v) fixative for 1 month and then examined via confocal microscopy and autofluorescence. Images were obtained with simultaneous excitation at 488 and 561 nm.

**Table 2 Significantly enriched gene ontologies associated with biological processes as deduced by DAVID from a list of differentially expressed genes identified by limma analysis**

Category	Term	Count	P value	Fold enrichment	BH	FDR
<b>Mxylem 1</b>						
GO:0010286	Heat acclimation	3	0.00	34.03	0.48	4.39
GO:0006952	Defence response	16	0.00	5.39	0.00	0.00
<b>Xylem 1</b>						
GO:0009813	Flavonoid biosynthetic process	15	0.00	18.17	0.00	0.00
GO:0009812	Flavonoid metabolic process	15	0.00	17.54	0.00	0.00
GO:0009699	Phenylpropanoid biosynthetic process	18	0.00	12.99	0.00	0.00
GO:0009698	Phenylpropanoid metabolic process	18	0.00	9.32	0.00	0.00
GO:0042398	Cellular amino acid derivative biosynthetic process	20	0.00	8.37	0.00	0.00
GO:0019438	Aromatic compound biosynthetic process	19	0.00	7.08	0.00	0.00
GO:0006575	Cellular amino acid derivative metabolic process	20	0.00	5.06	0.00	0.00
GO:0042542	Response to hydrogen peroxide	7	0.00	5.05	0.10	3.57
GO:0019748	Secondary metabolic process	21	0.00	4.88	0.00	0.00
GO:0006952	Defence response	15	0.00	3.36	0.01	0.20
<b>Bark 8</b>						
GO:0009853	Photorespiration	5	0.00	17.49	0.01	0.23
GO:0019253	Reductive pentose-phosphate cycle	4	0.00	15.64	0.05	2.73
GO:0015977	Carbon utilisation by fixation of carbon dioxide	4	0.00	13.99	0.07	3.78
GO:0019685	Photosynthesis, dark reaction	4	0.00	13.29	0.07	4.39
GO:0042744	Hydrogen peroxide catabolic process	9	0.00	8.31	0.00	0.02
GO:0009813	Flavonoid biosynthetic process	7	0.00	8.31	0.01	0.26
GO:0043094	Cellular metabolic compound salvage	5	0.00	8.11	0.07	4.47
GO:0070301	Cellular response to hydrogen peroxide	9	0.00	8.08	0.00	0.02
GO:0009812	Flavonoid metabolic process	7	0.00	8.02	0.01	0.31
GO:0042743	Hydrogen peroxide metabolic process	9	0.00	7.98	0.00	0.02
<b>Mxylem 8</b>						
GO:0009813	Flavonoid biosynthetic process	34	0.00	7.53	0.00	0.00
GO:0009812	Flavonoid metabolic process	35	0.00	7.48	0.00	0.00
GO:0009699	Phenylpropanoid biosynthetic process	40	0.00	5.27	0.00	0.00
GO:0009698	Phenylpropanoid metabolic process	45	0.00	4.26	0.00	0.00
GO:0009626	Plant-type hypersensitive response	8	0.00	4.13	0.10	3.62
GO:0034050	Host programmed cell death induced by symbiont	8	0.00	4.13	0.10	3.62
GO:0009739	Response to gibberellin stimulus	11	0.00	3.50	0.05	1.34
GO:0042398	Cellular amino acid derivative biosynthetic process	41	0.00	3.14	0.00	0.00
GO:0019438	Aromatic compound biosynthetic process	44	0.00	3.00	0.00	0.00
GO:0009415	Response to water	18	0.00	2.94	0.01	0.15
<b>Xylem 8</b>						
GO:0001906	Cell killing	6	0.00	9.31	0.01	0.38
GO:0031640	Killing of cells of another organism	6	0.00	9.31	0.01	0.38
GO:0009812	Flavonoid metabolic process	29	0.00	8.54	0.00	0.00
GO:0009813	Flavonoid biosynthetic process	28	0.00	8.54	0.00	0.00
GO:0009225	Nucleotide-sugar metabolic process	7	0.00	7.03	0.01	0.45
GO:0006032	Chitin catabolic process	9	0.00	6.68	0.00	0.05

**Table 2 Significantly enriched gene ontologies associated with biological processes as deduced by DAVID from a list of differentially expressed genes identified by limma analysis (Continued)**

GO ID	Gene Ontology	Count	Benjamini-Hochberg	FDR	Discovery Rate
GO:0016998	Cell wall macromolecule catabolic process	9	0.00	6.15	0.09
GO:0006026	Aminoglycan catabolic process	9	0.00	6.15	0.09
GO:0009699	Phenylpropanoid biosynthetic process	32	0.00	5.81	0.00
GO:0009698	Phenylpropanoid metabolic process	42	0.00	5.47	0.00

Mxylem, mucilaginous xylem; 1 and 8 refer to the number of weeks after application of Ethepon in lanolin; BH, Benjamini-Hochberg correction; FDR is false discovery rate.

## Results

### Transcriptomic analysis of ethephon induction in *Pinus radiata* saplings

*Pinus radiata* transcriptomes were analysed by heterologous hybridisation to the *Pinus* microarray with cRNA generated from different tissues extracted from ethephon-treated and control plants. The transcriptomes in bark, mucilaginous xylem and xylem 1 and 8 WAT with 0.2 g of lanolin containing 0 or 1 mg ethephon were compared by two-colour hybridisation. Limma analysis of the data revealed that the mucilaginous xylem had 66 and 23,084 significantly different signals (relative to the pertinent control) 1 and 8 WAT with ethephon, respectively; similarly, bark had 127 and 1761 (1 and 8 WAT) whilst xylem had 1,352 and 12,718 probes exhibiting significantly different signals (1 and 8 WAT).

The relationships between the differentially expressed probes among tissues and across time are shown in the Venn diagrams in Figure 2. At 1 WAT, the mucilaginous xylem and the bark displayed comparable numbers of up- and down-regulated genes, but only one of these was co-expressed. More genes in the xylem showed significant down-regulation due to ethephon but only a very small fraction were co-expressed in the mucilaginous xylem (12 or 3.9%) or bark (5 or 1.6%). Despite the very low numbers of differentially expressed genes in the bark at 1 WAT, at least 50% of these were still expressed in the bark at 8 WAT. A lower proportion (25-34%) of genes that were differentially expressed in the xylem at 1 WAT was also found seven weeks later.

Differentially expressed (fold change >2, adj-*p* < 0.05) genes were systematically mapped to the associated biological annotation (e.g., Gene Ontology Terms, KEGG

pathways) using DAVID, and the most over-represented (enriched) biological annotation were statistically highlighted. Enrichment analysis increases the likelihood of identifying the biological processes most relevant to the experiment (Huang et al. 2008). The Functional Annotation Table of Gene Ontologies with False Discovery Rates less than 0.1, generated through the DAVID portal, are shown in Table 2. Ethephon had no significant effect on the bark transcriptome one week after application. The xylem transcriptome had a greater response to ethephon than the bark transcriptome resulting in an increase in the activity of genes associated with cellular amino acid derivatives and phenylpropanoid biosynthesis in xylem at 1 WAT and 8 WAT. Ethephon initially depressed defence responses in the xylem but, after eight weeks, ethephon application activated a wide range of genes related to plant defence. These included genes involved in phenylpropanoid and flavonoid biosynthesis, chitin detection and degradation, oxidation-reduction, and toxin catabolism. Lignin metabolic processes were significantly down-regulated by ethephon treatment.


In the bark, genes associated with structural constituents of the cell wall (GO:0005199) were activated by ethephon. The down-regulation of light-responsive genes in the bark was expected, but the suppression of genes associated with abiotic stress was unexpected.

### Validation of microarray signals by quantitative RT-PCR

To test the validity of signal quantitation by the *Pinus* microarray, 60-mer probes were selected and corresponding transcripts in cDNA transcribed from the original total RNA extracts used for microarray hybridisation were quantified by RT-PCR. Five primer pairs designed to

**Table 3 Microarray probe targets for quantitative RT-PCR validation**

Key process	Gene	Probe ID	5' Primer	3'-Primer	PCR validation
Catechin Biosynthesis	Anthocyanidin reductase	Contig38855a	TGGCATCGACGAAATCTACG	TTTCGCGGATTCTACAGCGT	Yes
Isoflavonoid biosynthesis	Isoflavone reductase	Contig32761c	CCCACGTTGAATACACATCAGT	ACGGCTCAAATAACGCTCGACT	Yes
Mono terpenoids	Pinene synthase	Contig819c	CTAACTAGGTCTCTGATTT	GATATCTTACCCACAACATAGC	Yes
Unknown	Unknown	Contig24580b	AAATATAATCTCTGTCTCATCTGCG	AAGCGAATGGTTCATGGTC	Yes
Cell Cycle	Cyclophilin	Pta121982552a	AGAGTCGATCTACGGCGAGA	CTTTACAAAGTTCTCATCGGCA	Yes



amplify a 45–60 bp segment within each probe (Table 4) generated fold change estimates that were in general agreement with the direction (up- or down-regulation) of the raw signal data from the microarrays (Figure 3). The sixth probe used as a constitutive background control (contig 12730c) gave a consistent signal. Four primer pairs failed to amplify a corresponding target which is a consequence of the higher specificity required for PCR amplification which will not always be compatible for the detection of heterologous targets.

#### Anatomical changes after long-term ethephon exposure

Most of the visible response to ethephon was limited to the treated area on the bark. There was little effect on stem girth and appearance 1 cm above or below the lanolin band (Figure 4). After 8 weeks of treatment, the 1 mg dose induced an increase in girth (16%) relative to the girth immediately above and below the treated area. Increased stem diameter was evident in the ethephon-treated area but not in the xylem. Most of the increased growth was manifested in increased amounts of phloem and cortex in the bark that displayed deep fissures and cracks, and sap bleeding that turned the normally colourless lanolin white. Higher ethephon dosages (5 mg and 25 mg treatments) induced more resinification and generated greater levels of opacity and whitening of the lanolin due to reaction with bark exudates, bark splitting and necrosis and reduced stem growth (data not shown).

The ethephon-treated stems exhibited an abnormal layer of unligified tissue in the outer xylem, which is indicated by an asterisk in Figure 4B. The number and size of the resin canals were unaffected by ethephon treatment. After 1 month of fixation in FAA, unstained sections examined for autofluorescence indicated the presence of strongly red fluorescent extractives in both wood and bark from ethephon-treated plants (Figures 4C and 5). The red fluorescent parenchymatous cells, scattered throughout the outer xylem, are poorly lignified (Figure 5).

## Discussion

### Microarray construction and validation

Next Generation Sequencing (NGS) techniques can be used to produce gene libraries for the construction of microarrays (Hiremath et al. 2011), however generating numerous fully sequenced and annotated transcriptomes is expensive and time consuming. It is more efficient to capture a comprehensive picture of gene activity of an organism through the combining and normalisation of existing transcriptomes derived from various tissues sampled temporally and from a wide variety of natural and experimental conditions. Extensive sequence databases exist for *Pinus taeda* that were generated from diverse experimental conditions by many research groups; less extensive databases are also available for other *Pinus* species. Pre-existing public gene databases for *Pinus* species were used to construct the *Pinus* microarray described in this paper. The *Pinus* data were processed using CAP3 for reassembly of contigs, to remove redundancy and to generate consensus contigs (Huang and Madan 1999; Min et al. 2009; Feldmesser et al. 2013; Huang et al. 2012) for the All-*Pinus* database. *In silico* validation by BLASTN analysis of the All-*Pinus* database and the probes constructed from it revealed >99% identity of aligned sequences with reference to a set of publicly available *P. radiata* sequences. These *in silico* results were corroborated by the results obtained after *P. radiata* cRNA hybridisation.

### General observations on the transcriptomes induced by ethephon

Large numbers of genes were identified with highly significant levels (adjusted *P*-value < 0.01) of differential expression induced by ethephon treatment, and these were dependent on both tissue type and duration of ethephon exposure (Figure 2). Ethephon is a chemical that diffuses into the underlying plant tissues following application then generates the potent plant hormone ethylene in slightly acid or basic conditions in the cytoplasm (Lieberman 1979) where the effect of ethephon on endogenous ethylene biosynthesis is transient. This effect occurred in less than 6 days in the case of apple cuttings (Sun and Bassuk 1993) although the effect of ethephon on *P. radiata* after



**Table 4 Significantly suppressed gene ontologies associated with biological processes as deduced by DAVID from a list of differentially expressed genes identified by limma analysis**


Category	Term	Count	P value	Fold enrichment	Benjamini	FDR
<b>Bark 8</b>						
GO:0048444	Floral organ morphogenesis	4	0.00	15.78	0.07	2.50
GO:0030244	Cellulose biosynthetic process	7	0.00	9.56	0.01	0.10
GO:0015833	Peptide transport	6	0.00	8.52	0.04	0.88
GO:0006857	Oligopeptide transport	5	0.00	8.07	0.10	4.56
GO:0030243	Cellulose metabolic process	8	0.00	5.07	0.05	1.41
GO:0009699	Phenylpropanoid biosynthetic process	13	0.00	4.91	0.00	0.02
GO:0009250	Glucan biosynthetic process	8	0.00	4.58	0.07	2.58
GO:0006073	Cellular glucan metabolic process	13	0.00	4.35	0.00	0.06
GO:0009698	Phenylpropanoid metabolic process	16	0.00	4.34	0.00	0.01
GO:0009414	Response to water deprivation	8	0.00	4.12	0.10	4.75
<b>Mxylem 8</b>						
GO:0046274	Lignin catabolic process	17	0.00	6.25	0.00	0.00
GO:0046271	Phenylpropanoid catabolic process	17	0.00	6.25	0.00	0.00
GO:0048444	Floral organ morphogenesis	6	0.00	5.15	0.14	4.91
GO:0030244	Cellulose biosynthetic process	16	0.00	4.75	0.00	0.00
GO:0009808	Lignin metabolic process	31	0.00	4.35	0.00	0.00
GO:0051258	Protein polymerisation	25	0.00	4.29	0.00	0.00
GO:0042219	Cellular amino acid derivative catabolic process	19	0.00	3.96	0.00	0.00
GO:0007018	Microtubule-based movement	28	0.00	3.54	0.00	0.00
GO:0009698	Phenylpropanoid metabolic process	56	0.00	3.30	0.00	0.00
GO:0009809	Lignin biosynthetic process	13	0.00	3.14	0.03	0.68
<b>Xylem 8</b>						
GO:0046274	Lignin catabolic process	8	0.00	4.40	0.11	2.32
GO:0046271	Phenylpropanoid catabolic process	8	0.00	4.40	0.11	2.32
GO:0006857	Oligopeptide transport	8	0.00	4.20	0.14	3.14
GO:0009808	Lignin metabolic process	19	0.00	3.99	0.00	0.00
GO:0009698	Phenylpropanoid metabolic process	41	0.00	3.61	0.00	0.00
GO:0009809	Lignin biosynthetic process	10	0.00	3.61	0.11	1.96
GO:0009699	Phenylpropanoid biosynthetic process	29	0.00	3.56	0.00	0.00
GO:0009813	Flavonoid biosynthetic process	16	0.00	3.30	0.01	0.09
GO:0016137	Glycoside metabolic process	10	0.00	3.21	0.19	4.74
GO:0009812	Flavonoid metabolic process	16	0.00	3.18	0.01	0.15

Legend as in Table 2.

1 week was minimal. Once formed, however, ethylene triggered gene expression from each type of tissue at 8 WAT, and these were different in composition (Figure 2) and function (Table 2). Some (25-50%) of the genes expressed at 1 WAT were found to be differentially expressed at 8 WAT, indicating that these genes may form the core response of these tissues to ethephon. Amongst these were at least 8 gene ontology classes that showed bidirectional responses (both up- and down-regulated) to ethephon in


the xylem (Table 2). Overall, ethephon acted as a suppressor rather than as a promoter of gene transcription.


In *P. radiata* saplings, the mucilaginous xylem exists as a thin layer located between bark and xylem and includes the undifferentiated 1–2 cell thick cambium layer, expanding cells and a few early secondary walls classifiable as xylem but not phloem (Cato et al. 2006). This mucilaginous layer was practically absent in ethephon-treated stems by 8 WAT such that the recovered tissues contained



proportionally more fibrous material comprising young xylem. At 8 WAT, extensive contamination of the 'tissue scrapes' with more mature xylem tissues from ethephon-treated plants is the most probable reason for the unusually high number (>23,000) of probes showing significantly different signals when compared with corresponding control

plants from which 'tissue scrapes' uncontaminated by xylem were obtained. For this reason, the data from this treatment will not be discussed further, but serve as a reminder of the need for robust tissue typing and tissue collection if between-tissue comparisons are to be attempted. When comparing different studies it is important to understand the influence of the composition of the woody portion of the stem from which the RNA is extracted. This RNA from woody tissues is expected to comprise primarily of RNA from the outermost layer of woody differentiating xylem, since the final autolysis of cellular contents is slow in conifer tracheids (Bollhöner et al. 2012) but it will also contain RNA from rays and resin duct parenchyma, as the latter can live for more than 30 years in *Pinus* (Nakaba et al. 2008). The xylem transcriptomes obtained from the 2-year-old saplings used in this study likely reflect developmentally young, but woody, outer xylem seen immediately below the gelatinous 'developing xylem' of older trees. An expressed sequence tagged library has previously been generated from such older xylem tissues in 10-year-old *P. taeda* (Kirst et al. 2003). Differential gene expression among samples obtained from different organs (xylem, embryo, megagametophytes) have also been reported in *P. taeda* (Loopstra and Sederoff 1995).





**Figure 5 Comparison of outer stem anatomy between control (A) and 5 mg ethephon treatment (B).** Both Phloem (Ph) and Traumatic xylem (Tx) appear abnormal and contain increased extractives. The extractives are concentrated in poorly lignified parenchymatous cells in the xylem. Cortex (Cx) and Cambium (Ca) are as indicated. The images are at the same scale. Scale bar = 1 mm.

#### Ethephon induction of secondary metabolism

As expected, ethephon treatment triggered a chain of events that manifested as bark cracking and necrosis, resinification, and chlorosis, and led to significantly different transcriptomes (Table 2) in all sampled tissues. Ethephon increased the transcription of genes associated with secondary metabolism, specifically of phenylpropanoids and flavonoids in xylem tissues (Table 2). Some of these secondary metabolites may be responsible for the red-fluorescent extractives that developed in ethephon-treated xylem after fixation with FAA (Figure 4C). These extractives seem to be limited to the young outer xylem nearest to the mucilaginous xylem. This is consistent with the hypothesis that, in conifers, the ethylene signal is transported to the mucilaginous xylem which is subsequently reprogrammed to generate cells with altered physiology to produce metabolites that counter the stress factor identified by the signalling pathway (Hudgins and Franceschi 2004). Resinification was more pronounced at the higher 5 and 25 mg ethephon dosages (data not shown); transcriptomic analyses were not attempted for these treatments due to inconsistent RNA extraction possibly due to the adverse effect of high resin content.

#### Ethephon and plant defence

Ethylene is involved in the regulation of genes involved in resistance, defence and wounding response (Van Zhong and Burns 2003; Berrocal-Lobo and Molina 2004; Harfouche et al. 2006). Ethephon elicited greater transcript activity from genes implicated in defence against fungal and insect attack and oxidative stress (Table 2). Genes associated with binding (i.e. recognising) chitin, a

primary component of fungal cell walls and insect exoskeletons, and chitinases, enzymes designed to degrade chitins, were up-regulated in the xylem. Ethephon exposure also led to the up-regulation of genes related to oxidative stress; active oxygen species, such as the superoxide radical generated during pathogen attack and abiotic stress, are recognised by plants as a signal for triggering defence responses (Vranova et al. 2002).

#### Ethephon and anatomical structure of *P. radiata*

An increase in girth within the treated area (Figure 4) was similar to that reported for balsam fir *Abies balsamea* (L.) Mill. (Eklund and Little 1996), indicating that the effect of the ethylene generated from ethephon is limited to the area of application. The thickened cortex displayed deep fissures that did not extend to the xylem. Stereomicroscopic examination of the cut stem revealed that most of the increased growth was due to increased amounts of phloem and cortex rather than the woody portion of the stem. This confirms a previous report (Yamamoto and Kozlowski 1987a) where a 1% ethephon dose, which is approximately equal to the 1 mg ethephon treatment in this study, was shown to elicit the most bark growth. No significant increase in diameter in the xylem portion of the stem cross sections was observed (see Figure 4) and the increased growth on the xylem side of the mucilaginous xylem consisted of traumatic parenchyma rather than xylem tissue. There was no evidence of increased lignification or increased volume in the woody cross sections. In fact, stems exposed to 1 mg ethephon exhibited an abnormal layer of unlignified tissue in the outer xylem (asterisk in Figure 4B). This corroborates

the down-regulation of a cohort of genes identified via the microarray that are associated with lignin biosynthesis in the xylem which is consistent with a previous claim that ethephon reduced lignification of tracheids in compression wood that developed on the underside of tilted stems of *Pinus densiflora* (Yamamoto and Kozlowski 1987b).

#### Oleoresin and resin canals

In response to biotic stress, conifers secrete an oleoresin that is composed mainly of mono- and diterpenes (Schmidt et al. 2010). Oleoresin production may be associated with resin duct formation in the cambium zone and developing xylem in response to insect attack or experimentally following methyl jasmonate treatment (Zulak and Bohlmann 2010). In this study, applications of >1 mg of ethephon induced significant resin secretion although there were no clear differences in resin duct formation (Figure 4) over the duration of the experiment. The traumatic xylem that developed in stems treated with ethephon showed poorly lignified parenchymatous cells suffused with extractives (Figure 5). The lack of change in the size and number of the resin canals indicates that the effect of ethephon did not extend to the type of resin-duct formation that is associated with terpene synthesis over the time course of this experiment. By contrast, a similar treatment (1% ethephon in lanolin) had been found previously to induce significant resinification as well as terpene-duct formation in *P. halepensis* (Yamamoto and Kozlowski 1987a). Traumatic resin-duct formation after ethephon injection has also been observed in four-year-old Douglas-fir (*Pseudotsuga menziesii* (Mirb.) Franco) and giant redwood trees (*Sequoiadendron giganteum* Lindl. Buchholz) (Hudgins and Franceschi 2004). The higher levels of ethephon (5 and 25 mg) used in this study induced greater resin bleeding and bark splitting, and a reduction in stem girth and plant vitality leading to chlorosis.

#### Microarray and quantitative RT-PCR

Heterologous hybridisation of *Pinus radiata* derived probes to the *Pinus* microarray may produce spurious hybridisation results due to imperfect annealing that can lead to incorrect interpretations of gene expression levels; quantitative RT-PCR is therefore necessary to verify the microarray results. Quantitative real-time PCR is commonly used to validate gene expression results obtained from microarray analysis however there are factors inherent in these two methods that can lead to contradictory data (Morey et al. 2006). Microarrays are relatively insensitive in detecting <2-fold changes (Wang et al. 2006). For example, 28% of the rice genes selected for validation gave non-significant correlations between microarray and quantitative RT-PCR signals (Dubouzet et al. 2007), which was taken to be within normal expectations

for microarray experiments. In this study, five probes were identified with consistent fold-change values by quantitative RT-PCR that confirmed the direction of expression revealed by the microarray and three of the genes had almost identical fold change estimates. Four of the probes tested could not be amplified, indicating a difference in the ability to detect target transcripts between the heterologous microarray and the PCR platforms. A greater level of probe mismatching is tolerated within a microarray hybridisation (1–6 base differences over 60-mer probe can reduce signal intensity by at least 50% (LeProust 2008; Hughes et al. 2001)) compared with lower levels of mismatch that are permissible within a quantitative PCR experimental design. Mismatches may prevent successful RT-PCR validation with particular PCR primers or for the more dissimilar heterologous genes. In some cases, successful RT-PCR validation may be overcome through the fine tuning of PCR conditions including the re-design of an appropriate target within the microarray probe or target gene on which to base primer design so that it avoids areas of mismatch in the heterologous gene while still generating >45 bp fragments. Experimental factors such as using different reverse transcriptase-PCR kits or thermal cycling conditions may also need to be explored. It is important to note that a microarray is a tool to uncover genes of interest under particular experimental designs, and that those identified need to be confirmed using other approaches including quantitative RT-PCR. An important experimental consideration therefore is to ensure that sufficient material is archived to allow the required testing that may be required for validation.

#### Conclusion

It can be concluded that an annotated database of *Pinus* cDNA sequences can provide a sound basis for the design of a *Pinus* microarray to study gene expression in *P. radiata* by heterologous hybridisation. *In silico* comparisons indicated a very high similarity between publicly available transcripts derived from *P. radiata* and the microarray probes that were predominantly derived from *P. taeda*. Further genome sequencing initiatives within the Pinaceae have the potential to further expand the database upon which microarrays can be designed to interrogate transcriptome data across related species. The *Pinus* microarray was tested by heterologous hybridisation with cRNA produced from bark, mucilaginous xylem and xylem of *P. radiata* clonal saplings that were subjected to an ethephon induction trial. The microarray was able to highlight that ethephon initiated very different transcriptomes for all tissues sampled at 1 and 8 WAT and these could be associated with an expected cascade of physiological responses. As is common practice with microarray experiments validation of gene expression highlighted by the *Pinus* microarray via

quantitative RT-PCR is required; a caveat for validation of heterologous hybridisation is that primer design needs to overcome sequence differences that can exist between the heterologous test probes and the microarray 60-mer sequences. While not further tested here, the *Pinus* microarray developed may be useful for *P. taeda* and other closely related species in the Pinaceae since the majority (>98%) of sequences originate from *P. taeda*. Further development of the All-*Pinus* gene database is also valuable as more sequence data from related species in the Pinaceae becomes available.

#### Competing interests

The authors declare that they have no competing interests.

#### Authors' contributions

All authors contributed to this paper. JDG contributed initial idea and concept development, microarray design, experimental manipulations and manuscript writing; LD provided microscopy and interpretation; MAB and LM implemented microarray hybridisation and analysis; VL contributed database construction; and GLJ contributed to concept development and manuscript writing. All authors read and approved the final manuscript.

#### Acknowledgments

The authors wish to express their gratitude to the following Scion staff: Lucy Macdonald, Lisa Stanbra, Lorelle Phillips and, most specially, Heather Flint for their critical support in various technical aspects of this research. Drs. Tim Strabala and Armin Wagner provided invaluable insight and discussions that vastly improved the manuscript. Details of the microarray experiment have been submitted according to the specifications of the National Centre for Biotechnology Information (NCBI) Gene Expression Omnibus (GEO) and are available at <http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE42300>.

#### Author details

<sup>1</sup>Scion, 49 Sala St., Private Bag 3020, Rotorua, New Zealand. <sup>2</sup>Department of Biochemistry, University of Otago, Dunedin, New Zealand.

Received: 21 March 2014 Accepted: 3 September 2014

Published online: 17 October 2014

#### References

- Adomas, A, Heller, G, Li, G, Olson, A, Chu, T-M, Osborne, J, Craig, D, van Zyl, L, Wolfinger, R, Sederoff, R, Dean, RA, Stenlid, J, Finlay, R, & Asiegbu, FO. (2007). Transcript profiling of a conifer pathosystem: response of *Pinus sylvestris* root tissues to pathogen (*Heterobasidion annosum*) invasion. *Tree Physiology*, 27(10), 1441–1458.
- Altschul, SF, Madden, TL, Schäffer, AA, Zhang, J, Zhang, Z, Miller, W, & Lipman, DJ. (1997). Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Research*, 25(17), 3389–3402.
- Balk, P, Bronnum, P, Perks, M, Statten, E, van der Geest, L, & van Wordragen, M. (2007). Innovative Cold Tolerance Test for Conifer Seedlings. In In L. E. D. Riley, R. K.; Landis, T. D., tech. coords (Ed.), *National Proceedings: Forest and Conservation Nursery Associations-2006. RMRS-P-50* (Vol. RMRS-P-50, pp. 9–14). Fort Collins, CO: USDA Forest Service, Rocky Mountain Research Station.
- Benjamini, Y, & Hochberg, Y. (1995). Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society. Series B (Methodological)*, 57(1), 289–300.
- Berocal-Lobo, M, & Molina, A. (2004). Ethylene response factor 1 mediates Arabidopsis resistance to the soilborne fungus *Fusarium oxysporum*. *Molecular Plant-Microbe Interactions*, 17(7), 763–770.
- Bollhöner, B, Prestele, J, & Tuominen, H. (2012). Xylem cell death: emerging understanding of regulation and function. *Journal of Experimental Botany*, 63(3), 1081–1094.
- Brown, K, & Leopold, A. (1973). Ethylene and the regulation of growth in pine. *Canadian Journal of Forest Research*, 3(1), 143–145.
- Cato, S, McMillan, L, Donaldson, L, Richardson, T, Echt, C, & Gardner, R. (2006). Wood formation from the base to the crown in *Pinus radiata*: gradients of tracheid wall thickness, wood density, radial growth rate and gene expression. *Plant Molecular Biology*, 60(4), 565–581.
- Davey, M, Graham, N, Vanholme, B, Swennen, R, May, S, & Keulemans, J. (2009). Heterologous oligonucleotide microarrays for transcriptomics in a non-model species; a proof-of-concept study of drought stress in *Musa*. *BMC Genomics*, 10(1), 436.
- Dubouzet, JG, Ishihara, A, Matsuda, F, Miyagawa, H, Iwata, H, & Wakasa, K. (2007). Integrated metabolomic and transcriptomic analyses of high-tryptophan rice expressing a mutant anthranilate synthase alpha subunit. *Journal of Experimental Botany*, 58(12), 3309–3321.
- Eklund, L, & Little, CHA. (1996). Laterally applied Ethrel causes local increases in radial growth and indole-3-acetic acid concentration in *Abies balsamea* shoots. *Tree Physiology*, 16(5), 509–513.
- Feldmesser, E, Rosenwasser, S, Vardi, A, & Ben-Dor, S. (2013). Improving automated de-novo transcriptome definition in non-model organisms by integrating manually defined gene information. *EMBnet. Journal*, 19(A), 45–46.
- Hall, TA. (1999). BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nuclear Acids Symposium Series*, 41, 95–98.
- Harfouche, AL, Shivaji, R, Stocker, R, Williams, PW, & Luthe, DS. (2006). Ethylene signaling mediates a maize defense response to insect herbivory. *Molecular Plant Microbe Interactions: MPMI*, 19(2), 189–199.
- Heath, LS, Ramakrishnan, N, Sederoff, RR, Whetten, RW, Chevone, BI, Struble, CA, Jouenne, YV, Chen, D, van Zyl, L, & Grene, R. (2002). Studying the functional genomics of stress responses in loblolly pine with the Expresso microarray experiment management system. *Comparative and Functional Genomics*, 3(3), 226–243.
- Hiremath, PJ, Farmer, A, Cannon, SB, Woodward, J, Kudapa, H, Tuteja, R, Kumar, A, Bhanuprakash, A, Mulaosmanovic, B, Gujaria, N, Krishnamurthy, L, Gaur, PM, Kavikishor, PB, Shah, T, Srinivasan, R, Lohse, M, Xiao, Y, Town, CD, Cook, DR, May, GD, & Varshney, RK. (2011). Large-scale transcriptome analysis in chickpea (*Cicer arietinum* L.), an orphan legume crop of the semi-arid tropics of Asia and Africa. *Plant Biotechnology Journal*, 9(8), 922–931.
- Huang, X, & Madan, A. (1999). CAP3: A DNA sequence assembly program. *Genome Research*, 9(9), 868.
- Huang, DW, Sherman, BT, & Lempicki, RA. (2008). Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nature Protocols*, 4(1), 44–57.
- Huang, DW, Sherman, BT, & Lempicki, RA. (2009). Bioinformatics enrichment tools: paths toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Research*, 37(1), 1–13.
- Huang, Y-L, Fang, X-T, Lu, L, Yan, Y-B, Chen, S-F, Hu, L, Zhu, C-C, Ge, X-J, & Shi, S-H. (2012). Transcriptome analysis of an invasive weed *Mikania micrantha*. *Biologia Plantarum*, 56(1), 111–116.
- Hudgins, J, & Franceschi, V. (2004). Methyl jasmonate-induced ethylene production is responsible for conifer phloem defense responses and reprogramming of stem cambial zone for traumatic resin duct formation. *Plant Physiology*, 135(4), 2134.
- Hughes, T, Mao, M, Jones, A, Burchard, J, Marton, M, Shannon, K, Lefkowitz, SM, Ziman, M, Schelter, JM, Meyer, MR, Kobayashi, S, Davis, C, Dai, H, He, YD, Stephanians, SB, Cavet, G, Walker, WL, West, A, Coffey, E, Shoemaker, DD, Stoughton, R, Blanchard, AP, Friend, SH, & Linsley, PS. (2001). Expression profiling using microarrays fabricated by an ink-jet oligonucleotide synthesizer. *Nature Biotechnology*, 19(4), 342–347.
- Kirst, M, Johnson, A, Baucom, C, Ulrich, E, Hubbard, K, Staggs, R, Paule, C, Retzel, E, Whetten, R, & Sederoff, R. (2003). Apparent homology of expressed genes from wood-forming tissues of loblolly pine (*Pinus taeda* L.) with *Arabidopsis thaliana*. *Proceedings of the National Academy of Sciences*, 100(12), 7383–7388.
- LeProust, E. (2008). Agilent's Microarray Platform: How High-Fidelity DNA Synthesis Maximizes the Dynamic Range of Gene Expression Measurements Agilent's Application Notes (p. 12), Agilent Technologies ([www.agilent.com/chem/dna](http://www.agilent.com/chem/dna)).
- Li, X, Wu, HX, & Southerton, SG. (2012). Identification of putative candidate genes for juvenile wood density in *Pinus radiata*. *Tree Physiology*, 32(8), 1046–1057.
- Lieberman, M. (1979). Biosynthesis and action of ethylene. *Annual Review of Plant Physiology*, 30(1), 533–591.
- Loopstra, CA, & Sederoff, RR. (1995). Xylem-specific gene expression in loblolly pine. *Plant Molecular Biology*, 27(2), 277–291.
- Lorenz, WW, Alba, R, Yu, YS, Bordeaux, J, Simoes, M, & Dean, J. (2011). Microarray analysis and scale-free gene networks identify candidate regulators in drought-stressed roots of loblolly pine (*P. taeda* L.). *BMC Genomics*, 12(1), 264.

- Lorenz, WW, Ayyampalayam, S, Bordeaux, JM, Howe, GT, Jermstad, KD, Neale, DB, Rogers, DL, & Dean, JFD. (2012). Conifer DBMAGIC: a database housing multiple *de novo* transcriptome assemblies for 12 diverse conifer species. *Tree Genetics Genomics*, *8*, 1477–1485.
- Min, XJ, Butler, G, Storms, R, & Tsang, A. (2009). Comparative Assessment of DNA Assemblers for Assembling Expressed Sequence Tags. In *Bioinformatics (OCCBIO'09. Ohio Collaborative Conference on, 2009*, pp. 79–82). IEEE. doi:10.1109/OCCBIO.2009.19.
- Morey, JS, Ryan, JC, & Van Dolah, FM. (2006). Microarray validation: factors influencing correlation between oligonucleotide microarrays and real-time PCR. *Biological Procedures Online*, *8*(1), 175–193.
- Myburg, H, Morse, A, Amerson, H, Kubisiak, T, Huber, D, Osborne, J, Osborne, JA, Garcia, SA, Nelson, CD, Davis, JM, Covert, SF, & van Zyl, LM. (2006). Differential gene expression in loblolly pine (*Pinus taeda* L.) challenged with the fusiform rust fungus, *Cronartium quercuum* f. sp. *fusiforme*. *Physiological and Molecular Plant Pathology*, *68*(1–3), 79–91.
- Nakaba, S, Kubo, T, & Funada, R. (2008). Differences in patterns of cell death between ray parenchyma cells and ray tracheids in the conifers *Pinus densiflora* and *Pinus rigida*. *Trees-Structure and Function*, *22*(5), 623–630.
- New Zealand Forest Owners Association. (2011). New Zealand Plantation Forest Industry: Facts and Figures. In N. Z. F. O. Association (Ed.), (Vol. 2010/11, p. 4). New Zealand: New Zealand Forest Owners Association.
- Perdiguero, P, Collada, C, Barbero, MC, Casado, GG, Cervera, MT, & Soto, Á. (2012). Identification of water stress genes in *Pinus pinaster* Ait. by controlled progressive stress and suppression-subtractive hybridization. *Plant Physiology and Biochemistry*, *50*(1), 44–53.
- Pfaffl, MW. (2004). Quantification strategies in real-time PCR. *AZ of Quantitative PCR*, *1*, 89–113.
- Reinders, J, Vivier, CD, Theiler, G, Chollet, D, Descombes, P, & Paszkowski, J. (2008). Genome-wide, high-resolution DNA methylation profiling using bisulfite-mediated cytosine conversion. *Genome Research*, *18*(3), 469–476.
- Schmidt, A, Wächtler, B, Temp, U, Krekling, T, Séguin, A, & Gershenzon, J. (2010). A bifunctional geranyl and geranylgeranyl diphosphate synthase is involved in terpene oleoresin formation in *Picea abies*. *Plant Physiology*, *152*(2), 639–655.
- Schnurr, JP, Cheng, Z, & Boe, AA. (1996). Effects of plant growth regulators on sturdiness of Jack pine seedlings. *Journal of Environmental Horticulture*, *14*, 228–230.
- Smyth, GK. (2005). Limma: Linear Models for Microarray Data. In R Gentleman, VJ Carey, W Huber, RA Irizarry, & S Dudoit (Eds.), *Bioinformatics and Computational Biology Solutions Using R and Bioconductor* (Bioinformatics and Computational Biology Solutions Using R and Bioconductor, pp. 397–420). New York: Springer Science + Business Media New York.
- Sun, WQ, & Bassuk, NL. (1993). Auxin-induced Ethylene Synthesis during Rooting and Inhibition of Budbreak of Royalty/Rose Cuttings. *Journal of the American Society for Horticultural Science*, *118*(5), 638–643.
- Van Zhong, G, & Burns, JK. (2003). Profiling ethylene-regulated gene expression in *Arabidopsis thaliana* by microarray analysis. *Plant Molecular Biology*, *53*(1), 117–131.
- Van Zyl, L, Von Arnold, S, Bozhkov, P, Chen, Y, Egertsdotter, U, MacKay, J, Sederoff, RR, Shen, J, Zelena, L, & Clapham, DH. (2002). Heterologous array analysis in Pinaceae: hybridization of *Pinus taeda* cDNA arrays with cDNA from needles and embryogenic cultures of *P. taeda*, *P. sylvestris* or *Picea abies*. *Comparative and Functional Genomics*, *3*(4), 306–318.
- Vranova, E, Inzé, D, & Van Breusegem, F. (2002). Signal transduction during oxidative stress. *Journal of Experimental Botany*, *53*(372), 1227–1236.
- Wang, Y, Barbacioru, C, Hyland, F, Xiao, W, Hunkapiller, KL, Blake, J, Chan, F, Gonzalez, C, Zhang, L, & Samaha, RR. (2006). Large scale real-time PCR validation on gene expression measurements from two commercial long-oligonucleotide microarrays. *BMC Genomics*, *7*(1), 59.
- Warner, H, & Leopold, A. (1969). Ethylene evolution from 2-chloroethylphosphonic acid. *Plant Physiology*, *44*(1), 156.
- Yamamoto, F, & Kozlowski, T. (1987a). Effect of ethrel on growth and stem anatomy of *Pinus halepensis* seedlings. *IAWA Bulletin*, *8*(1), 11–19.
- Yamamoto, F, & Kozlowski, T. (1987b). Effects of flooding, tilting of stems, and ethrel application on growth, stem anatomy and ethylene production of *Pinus densiflora* seedlings. *Journal of Experimental Botany*, *38*(2), 293–310.
- Yang, S-H, van Zyl, L, No, E-G, & Loopstra, CA. (2004). Microarray analysis of genes preferentially expressed in differentiating xylem of loblolly pine (*Pinus taeda*). *Plant Science*, *166*(5), 1185–1195.
- Zulak, KG, & Bohlmann, J. (2010). Terpenoid biosynthesis and specialized vascular cells of conifer defense. *Journal of Integrative Plant Biology*, *52*(1), 86–97.

doi:10.1186/s40490-014-0021-8

Cite this article as: Dubouzet et al.: Heterologous hybridisation to a *Pinus* microarray: profiling of gene expression in *Pinus radiata* saplings exposed to ethephon. *New Zealand Journal of Forestry Science* 2014 **44**:21.

Submit your manuscript to a SpringerOpen<sup>®</sup> journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► [springeropen.com](http://springeropen.com)